

# THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

*June 16, 2005*

**THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.**

**APPLICATION NUMBER: 60/577,361**

**FILING DATE: *June 04, 2004***

**RELATED PCT APPLICATION NUMBER: *PCT/US05/17167***



Certified by

Under Secretary of Commerce  
for Intellectual Property  
and Director of the United States  
Patent and Trademark Office

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Provisional Application of  
DAVID W. OLD, et al.

For: 5-THIO-PIPERIDINYL PROSTAGLANDIN E  
ANALOGS AS THERAPEUTIC AGENTS

**PROVISIONAL APPLICATION FOR PATENT COVER SHEET**

Commissioner for Patents  
P.O. Box 1450  
Alexandria, VA 22313-1450

Dear Sir:

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT  
under 37 CFR 1.53(c).

1. The enclosed PROVISIONAL APPLICATION includes the following:
  - a. Specification (32 pgs);
  - b. Claims (5 pgs);
  - c. Abstract (1 pg);
  - d. Drawings (2 sheets).

2. Inventors:

Given Name	Residence
DAVID W. OLD	13771 TYPEE WAY IRVINE, CA 92620
DANNY T. DINH	11531 COLLEGE AVENUE GARDEN GROVE, CA 92840

3. Fee Payment

Please charge to Deposit Account No. 01-0885 for the filing fee of \$160.00, as well as for any discrepancies that may occur.

4. Postcard Receipt

Applicant submits herewith a Return/Stamped Postcard.

Respectfully submitted,

Date: June 3, 2004

By: 

Brent A. Johnson, Ph.D.  
Patent Agent  
Registration No. 51,851  
Tel: 714/246-4348  
Fax: 714/246-4249


Please address all future correspondence to:

Brent A. Johnson, Ph.D.  
ALLERGAN, INC.  
2525 Dupont Drive, T2-7H  
Irvine, CA 92612  
Tel: 714/246-4348  
Fax: 714/246-4249

**CERTIFICATE OF EXPRESS MAIL UNDER 37 C.F.R. §1.10**

I hereby certify that this Provisional Patent Application and the documents referred to as enclosed herein are being deposited with the United States Postal Service on JUNE 3, 2004 in an envelope as "Express Mail Post Office To Addressee" mailing label number EV193717023US with sufficient postage for Express Mail addressed to Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

Susan Bartholomew  
Name of person mailing paper

  
Signature of person mailing paper

Date: JUNE 3, 2004

5

**5-THIO-PIPERDINYL PROSTAGLANDIN E ANALOGS****By Inventors****David W. Old and Danny T. Dinh****FIELD OF THE INVENTION**

10

This invention relates to compounds which are useful as therapeutic agents. Among other potential uses, these compounds are believed to have properties which are characteristic of prostaglandins.

15

**BACKGROUND OF THE INVENTION****Description of Related Art**

Ocular hypotensive agents are useful in the treatment of a number of various ocular hypertensive conditions, such as post-surgical and post-laser trabeculectomy ocular hypertensive episodes, glaucoma, and as presurgical adjuncts.

Glaucoma is a disease of the eye characterized by increased intraocular pressure. On the basis of its etiology, glaucoma has been classified as primary or secondary. For example, primary glaucoma in adults (congenital glaucoma) may be either open-angle or acute or chronic angle-closure. Secondary glaucoma results from pre-existing ocular diseases such as uveitis, intraocular tumor or an enlarged cataract.


The underlying causes of primary glaucoma are not yet known. The increased intraocular tension is due to the obstruction of aqueous humor outflow. In chronic open-angle glaucoma, the anterior chamber and its anatomic structures appear normal, but drainage of the aqueous humor is impeded. In acute or chronic angle-closure glaucoma, the anterior chamber is shallow, the filtration angle is narrowed, and the iris may obstruct the trabecular meshwork at the entrance of the canal of Schlemm. Dilation of the pupil may push the root of the iris forward against the angle, and may produce pupillary block and thus

5 precipitate an acute attack. Eyes with narrow anterior chamber angles are predisposed to acute angle-closure glaucoma attacks of various degrees of severity.

Secondary glaucoma is caused by any interference with the flow of aqueous humor from the posterior chamber into the anterior chamber and subsequently, into the canal of Schlemm. Inflammatory disease of the anterior segment may prevent aqueous escape by causing complete posterior synechia in iris bombe, and may plug the drainage channel with exudates. Other common causes are intraocular tumors, enlarged cataracts, central retinal vein occlusion, trauma to the eye, operative procedures and intraocular hemorrhage.

15 Considering all types together, glaucoma occurs in about 2% of all persons over the age of 40 and may be asymptotic for years before progressing to rapid loss of vision. In cases where surgery is not indicated, topical  $\beta$ -adrenoreceptor antagonists have traditionally been the drugs of choice for treating glaucoma.

20 Certain eicosanoids and their derivatives have been reported to possess ocular hypotensive activity, and have been recommended for use in glaucoma management. Eicosanoids and derivatives include numerous biologically important compounds such as prostaglandins and their derivatives. Prostaglandins can be described as derivatives of prostanoic acid which have the following structural formula:



Various types of prostaglandins are known, depending on the structure and substituents carried on the alicyclic ring of the prostanoic acid skeleton.

30 Further classification is based on the number of unsaturated bonds in the side

5 chain indicated by numerical subscripts after the generic type of prostaglandin [e.g. prostaglandin E<sub>1</sub> (PGE<sub>1</sub>), prostaglandin E<sub>2</sub> (PGE<sub>2</sub>)], and on the configuration of the substituents on the alicyclic ring indicated by  $\alpha$  or  $\beta$  [e.g. prostaglandin F<sub>2</sub> $\alpha$  (PGF<sub>2</sub> $\beta$ )].

Prostaglandins were earlier regarded as potent ocular hypertensives,  
10 however, evidence accumulated in the last decade shows that some prostaglandins are highly effective ocular hypotensive agents, and are ideally suited for the long-term medical management of glaucoma (see, for example, Bito, L.Z. Biological Protection with Prostaglandins, Cohen, M.M., ed., Boca Raton, Fla, CRC Press Inc., 1985, pp. 231-252; and Bito, L.Z., Applied  
15 Pharmacology in the Medical Treatment of Glaucomas Drance, S.M. and Neufeld, A.H. eds., New York, Grune & Stratton, 1984, pp. 477-505. Such prostaglandins include PGF<sub>2</sub> $\alpha$ , PGF<sub>1</sub> $\alpha$ , PGE<sub>2</sub>, and certain lipid-soluble esters, such as C<sub>1</sub> to C<sub>2</sub> alkyl esters, e.g. 1-isopropyl ester, of such compounds.

Although the precise mechanism is not yet known experimental results  
20 indicate that the prostaglandin-induced reduction in intraocular pressure results from increased uveoscleral outflow [Nilsson et. al., Invest. Ophthalmol. Vis. Sci. (suppl), 284 (1987)].

The isopropyl ester of PGF<sub>2</sub> $\alpha$  has been shown to have significantly greater hypotensive potency than the parent compound, presumably as a result of  
25 its more effective penetration through the cornea. In 1987, this compound was described as "the most potent ocular hypotensive agent ever reported" [see, for example, Bito, L.Z., Arch. Ophthalmol. 105, 1036 (1987), and Siebold et al., Prodrug 5 3 (1989)].

Whereas prostaglandins appear to be devoid of significant intraocular side  
30 effects, ocular surface (conjunctival) hyperemia and foreign-body sensation have been consistently associated with the topical ocular use of such compounds, in particular PGF<sub>2</sub> $\alpha$  and its prodrugs, e.g., its 1-isopropyl ester, in humans. The clinical potentials of prostaglandins in the management of conditions associated with increased ocular pressure, e.g. glaucoma are greatly limited by these side  
35 effects.

5           In a series of United States patents assigned to Allergan, Inc.  
prostaglandin esters with increased ocular hypotensive activity accompanied with  
no or substantially reduced side-effects are disclosed. Some representative  
examples are U.S. Patent 5,446,041, U.S. Patent 4,994,274, U.S. Patent  
5,028,624 and U.S. Patent 5,034,413 all of which are hereby expressly  
10   incorporated by reference.

US Patent No. 5,688, 819, commonly assigned to Allergan, Inc., and  
incorporated herein by reference discloses compounds known as prostamides.  
Prostamides are distinguished from prostaglandins in that the oxygen which is  
bonded to carbonyl group is replaced by a nitrogen bearing substituent. Those  
15   skilled in the art will readily recognize that this replacement significantly alters  
several electronic and steric properties of an important structural feature in the  
biological molecule. Significantly, it is commonly believed in the art that  
resonance between the nitrogen lone pair and the carbonyl  $\pi$ -bond is  
significantly greater than resonance between the carbonyl group and an oxygen  
20   lone pair in a carboxylic ester or a carboxylic acid. This belief is supported by  
the well established experimental observation that the nitrogen atom in an  
amide is planar, as opposed to the pyramidal geometry of an amine. Thus, the  
commonly accepted belief in the art is that the nitrogen atom of an amine is  $sp^3$   
hybridized, while nitrogen atom of an amide is  $sp^2$  hybridized, with the bonded  
25   electrons occupying the  $sp^2$  hybrid orbitals and the nonbonded electron pair  
occupying a p orbital to allow for conjugation with the carbonyl  $\pi$  system. By  
contrast, the hybridization, bonding, and geometry of the electrons of the  
oxygen atom in water and alcohols are very similar to those of carboxylic acids  
or carboxylic esters.

30           The increased resonance between the nitrogen and the carbonyl group in  
the amide confers several unique properties to the molecule. First, it is well  
known in the art that hydrolysis of amides is at least two orders of magnitude  
slower than the hydrolysis of esters (see, for example, Francis A. Carey,  
Organic Chemistry, New York: McGraw-Hill Book Company, 1987, p. 779).  
35   Thus, hydrolysis of amides in vivo is slowed to such an extent that a prostamide  
cannot be considered to be a prodrug of a prostaglandin. Second, the increased


5 resonance significantly increases the barrier to rotation about the nitrogen-carbonyl sigma bond relative to the analogous rotational barrier associated with esters and carboxylic acids. Thus, a prostamide has a sterically significant, stable, rigid group replacing the oxygen atom of the prostaglandin. This significant steric difference will have a significant effect in binding to a number  
10 of receptor sites since geometry is important for many receptor sites. Since the carboxylic acid group of a prostaglandin is a polar, ionizable, group, with four potential hydrogen bond receiving electron pairs, and in the case of the protonated acid, one potential hydrogen bond donor, it is reasonable for a person of ordinary skill in the art to believe that this functional group will be  
15 important to the binding of the molecule to a number of receptors. It follows that changing the resonance properties, the hybridization of the bonding and nonbonding electrons, the geometry of the nitrogen atom, the number of available hydrogen bonding sites, and the electronegativity of the of the nitrogen relative to oxygen, will confer significantly different biological  
20 properties to prostamides relative to prostaglandins.

Recently, it is becoming more commonly accepted in the art that amides have distinct properties over carboxylic acids. For example, it has been shown that anandamide, a common amide of arachidonic acid, has significant biological activity that arachidonic acid does not. Other work has also been  
25 done to show that amides have distinct activity as compared to carboxylic acid, which has caused some in the field to classify fatty acid amides as "a new family of biologically active lipids" (Bezuglov, et. al., "Synthesis and Biological Evaluation of Novel Amides of Polyunsaturated Fatty Acids with Dopamine", *Bioorganic & Medicinal Chemistry Letters* 11 (2001), 447-449).

30 It has been shown that prostamides can have pronounced effects on smooth muscle and are potent ocular hypotensive agents. Additionally, prostamides may cause significantly lower ocular surface hyperemia than prostaglandins. One prostamide exemplary of the these effects is bimatoprost, which is marketed by Allergan, Inc. under the trade name Lumigan®, which has  
35 the structure shown below.



6




5

Although prostamide compounds have activity which is distinct from prostaglandins, they have many similar structural features. While not intending to be bound in any way by theory, it is believed that the structural similarity arises because prostamides are biosynthesized from N-arachidonyl ethanolamide whereas prostaglandins are biosynthesized from the structurally related arachidonic acid. Thus, they have similar structural traits, but play physiologically distinct roles due to the unique differences between the amide and the acid or ester functional groups highlighted previously. For example, it is believed that the two classes of compounds are active at distinct receptors. Thus, it is believed that the prostamide and prostaglandin receptors recognize a similar geometry in terms of the basic ring and  $\alpha$ - and  $\omega$ - chain structure, or analogs thereof, but selectively distinguish between prostaglandin and prostamide compounds based upon the nitrogen or oxygen substitution at the carbonyl group.

### **BRIEF DESCRIPTION OF THE INVENTION**

A compound comprising



5


or a pharmaceutically acceptable salt or a prodrug thereof, is disclosed herein, wherein a dashed line indicates the presence or absence of a bond, and an (H) represents a hydrogen atom which is present if required by said bond;

Y is selected from the group consisting of  $\text{CO}_2\text{H}$ ,  $\text{CONMe}_2$ ,  $\text{CONHMe}$ ,

10  $\text{CONHEt}$ ,  $\text{CON}(\text{OCH}_3)\text{CH}_3$ ,  $\text{CONH}_2$ ,  $\text{CON}(\text{CH}_2\text{CH}_2\text{OH})_2$ ,

$\text{CONH}(\text{CH}_2\text{CH}_2\text{OH})$ ,  $\text{CH}_2\text{OH}$ ,  $\text{P}(\text{O})(\text{OH})_2$ ,  $\text{CONHSO}_2\text{CH}_3$ ,  $\text{SO}_2\text{NH}_2$ ,

$\text{SO}_2\text{N}(\text{CH}_3)_2$ ,  $\text{SO}_2\text{NH}(\text{CH}_3)$ ,




, and

; and

R is selected from the group consisting of  $\text{C}_1$ - $\text{C}_4$  alkyl,  $\text{C}_1$ - $\text{C}_4$  alkoxy, halogen,

15  $\text{CO}_2\text{H}$ ,  $\text{OH}$ ,  $\text{COH}$ ,  $\text{COCH}_3$ ,  $\text{COCF}_3$ ,  $\text{NO}_2$ ,  $\text{CN}$ , and  $\text{CF}_3$ .

A compound having an  $\omega$  chain comprising



or a derivative thereof, is disclosed herein,

wherein a dashed line indicates the presence or absence of a bond, and an (H)

20 represents a hydrogen atom which is present if required by said bond;

- 5 wherein said derivative has a structure as shown above except that an alteration is made to said structure, wherein an alteration consists of
- a. adding, removing, or substituting a non-hydrogen atom of the  $\omega$  chain;
  - b. converting a  $\text{CO}_2\text{H}$  to a moiety selected from the group  
10 consisting of  $\text{CONMe}_2$ ,  $\text{CONHMe}$ ,  $\text{CONHEt}$ ,  $\text{CON}(\text{OCH}_3)\text{CH}_3$ ,  $\text{CONH}_2$ ,  $\text{CON}(\text{CH}_2\text{CH}_2\text{OH})_2$ ,  $\text{CONH}(\text{CH}_2\text{CH}_2\text{OH})$ ,  $\text{CH}_2\text{OH}$ ,  $\text{P}(\text{O})(\text{OH})_2$ ,  $\text{CONHSO}_2\text{CH}_3$ ,  $\text{SO}_2\text{NH}_2$ ,  $\text{SO}_2\text{N}(\text{CH}_3)_2$ ,  $\text{SO}_2\text{NH}(\text{CH}_3)$ ,
- The image shows two chemical structures. The first structure on the left is N-(cyclopropyl)acetamide, which consists of an acetamide group (a carbonyl group bonded to a methyl group and an NH group) where the hydrogen of the NH group is replaced by a cyclopropyl ring. The second structure on the right is 1-methyl-1H-1,2,4-triazole, which is a five-membered aromatic ring with two nitrogen atoms at positions 1 and 4, and a methyl group attached to the carbon at position 5. The two structures are separated by a comma and the word 'and'.
- c. converting a phenyl moiety to a pyridinyl, furyl, thienyl, or *n*-butyl moiety, or
  - d. adding a substituent comprising from 1 to 3 non-hydrogen atoms to a phenyl moiety;





- or a pharmaceutically acceptable salt or a prodrug thereof.
- 20 Methods of treating certain conditions or diseases, and compositions and medicaments related thereto are also contemplated.

### **BRIEF DESCRIPTION OF THE DRAWING FIGURES**

- 25 Figures 1 and 2 illustrate one method of preparing the compounds disclosed herein.

### **DETAILED DESCRIPTION OF THE INVENTION**

- 30 In the structures depicted herein, a dashed line indicates the presence or absence of a bond. Thus, while not intending to be limiting, the compounds shown below are possible.







5

Pharmaceutically acceptable salts or prodrugs of these compounds are also considered to be useful.

Additionally, the following compounds or derivatives thereof, or pharmaceutically acceptable salts or prodrugs of these compounds or derivatives are contemplated.

10



15

The phrase "an (H) represents a hydrogen atom which is present if required by said bond" is intended to mean that in the case that a bond indicated by a dashed line is not present, the hydrogen will be present to complete a C-OH moiety, as in some of the structures above. Alternatively if a dashed line indicates a bond which is part of a C=O moiety, no hydrogen is present.


A person of ordinary skill in the art understands the meaning of the stereochemistry associated with the hatched wedge/solid wedge structural

5 features. For example, an introductory organic chemistry textbook (Francis A. Carey, Organic Chemistry, New York: McGraw-Hill Book Company 1987, p. 63) states "a wedge indicates a bond coming from the plane of the paper toward the viewer" and the hatched wedge, indicated as a "dashed line", "represents a bond receding from the viewer."

10 "C<sub>1</sub>-C<sub>4</sub> alkyl" refers to any hydrocarbon having 1-4 carbon atoms and only single bonds, whether linear, branched, or cyclic, or a combination thereof. Thus, while not intending to be limiting, methyl, ethyl, propyl, isopropyl, *n*-butyl, isobutyl, *sec*-butyl, *tert*-butyl, cyclopropyl, cyclobutyl, methylcyclopropyl, and the like are "C<sub>1</sub>-C<sub>4</sub>" alkyl.

15 "C<sub>1</sub>-C<sub>4</sub> alkoxy" refers to moiety having O directly attached to the remaining part of the molecule and to a C<sub>1</sub>-C<sub>4</sub> alkyl. Thus, while not intending to be limiting, -O-methyl, -O-ethyl, -O-propyl, -O-isopropyl, -O-*n*-butyl, -O-isobutyl, -O-*sec*-butyl, -O-*tert*-butyl, -O-cyclopropyl, -O-cyclobutyl, -O-methylcyclopropyl, and the like are "C<sub>1</sub>-C<sub>4</sub>" alkoxy.

20 While not intending to limit the scope of the invention in any way, some compounds comprise




or a pharmaceutically acceptable salt or a prodrug thereof.

25 A "pharmaceutically acceptable salt" is any salt that retains the activity of the parent compound and does not impart any additional deleterious or untoward effects on the subject to which it is administered and in the context in which it is administered compared to the parent compound. A pharmaceutically acceptable salt also refers to any salt which may form in vivo as a result of administration of an acid, another salt, or a prodrug which is converted into an  
30 acid or salt.

5           Pharmaceutically acceptable salts of acidic functional groups may be derived from organic or inorganic bases. The salt may comprise a mono or polyvalent ion. Of particular interest are the inorganic ions, lithium, sodium, potassium, calcium, and magnesium. Organic salts may be made with amines, particularly ammonium salts such as mono-, di- and trialkyl amines or ethanol  
10 amines. Salts may also be formed with caffeine, tromethamine and similar molecules. Hydrochloric acid or some other pharmaceutically acceptable acid may form a salt with a compound that includes a basic group, such as an amine or a pyridine ring.


          A "prodrug" is a compound which is converted to a therapeutically  
15 active compound after administration, and the term should be interpreted as broadly herein as is generally understood in the art. While not intending to limit the scope of the invention, conversion may occur by hydrolysis of an ester group or some other biologically labile group. Ester prodrugs of the compounds disclosed herein are specifically contemplated. An ester may be derived from a  
20 carboxylic acid of C1 (i.e. the terminal carboxylic acid of a natural prostaglandin), or an ester may be derived from a carboxylic acid functional group on another part of the molecule, such as on a phenyl ring. While not intending to be limiting, an ester may be an alkyl ester, an aryl ester, or a heteroaryl ester. The term alkyl has the meaning generally understood by those  
25 skilled in the art and refers to linear, branched, or cyclic alkyl moieties. C<sub>1-6</sub> alkyl esters are particularly useful, where alkyl part of the ester has from 1 to 6 carbon atoms and includes, but is not limited to, methyl, ethyl, propyl, isopropyl, *n*-butyl, *sec*-butyl, *iso*-butyl, *t*-butyl, pentyl isomers, hexyl isomers, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and combinations thereof  
30 having from 1-6 carbon atoms, etc.

          While not intending to be limiting, one example of a prodrug consists of




5

The tetrazole group,



has two tautomeric forms, which can rapidly interconvert in aqueous or biological media, and are thus equivalent to one another. The tautomer of the tetrazole shown above is shown below.

10




For the purposes disclosed herein, all tautomeric forms should be considered equivalent in every way.

In making reference to a derivative and alterations to a structure, it should be emphasized that making alterations and forming derivatives is strictly a mental exercise used to define a set of chemical compounds, and has nothing to do with whether said alteration can actually be carried out in the laboratory, or whether a derivative can be prepared by an alteration described. However, whether the derivative can be prepared via any designated alteration or not, the differences between the derivatives and the aforementioned structure are such that a person of ordinary skill in the art could prepare the derivatives disclosed herein using routine methods known in the art without undue experimentation.

15

20





5

The  $\omega$  chain is the group circled in the labeled structure above.

Changes to the structure can take several forms, if a non-hydrogen atom is added, the structure is changed by adding the atom, and any required hydrogen atoms, but leaving the remaining non-hydrogen atoms unchanged, such as in the two examples shown below, with the added atoms in bold type.



10



Pharmaceutically acceptable salts, tetrazoles, and prodrugs of these compounds are also contemplated.

15

If a non-hydrogen atom is removed, the structure is changed by removing the atom, and any required hydrogen atoms, but leaving the remaining non-hydrogen atoms unchanged, such as in the two examples shown below, with the previous location of the missing atoms indicated by arrows.





20

Pharmaceutically acceptable salts, tetrazoles, and prodrugs of these compounds are also contemplated.




- 5 If a non-hydrogen atom is substituted, the non-hydrogen atom is replaced by a different non-hydrogen atom, with any necessary adjustment made to the number hydrogen atoms, such as in the two examples shown below, with the substituted atoms in bold type.




- 10 Pharmaceutically acceptable salts, tetrazoles, and prodrugs of these compounds are also contemplated.

Another alteration includes converting a  $\text{CO}_2\text{H}$  to a moiety selected from the group consisting of  $\text{CONMe}_2$ ,  $\text{CONHMe}$ ,  $\text{CONHEt}$ ,  $\text{CON}(\text{OCH}_3)\text{CH}_3$ ,  $\text{CONH}_2$ ,  $\text{CON}(\text{CH}_2\text{CH}_2\text{OH})_2$ ,  $\text{CONH}(\text{CH}_2\text{CH}_2\text{OH})$ ,  $\text{CH}_2\text{OH}$ ,  $\text{P}(\text{O})(\text{OH})_2$ ,





- 15  $\text{CONHSO}_2\text{CH}_3$ ,  $\text{SO}_2\text{NH}_2$ ,  $\text{SO}_2\text{N}(\text{CH}_3)_2$ ,  $\text{SO}_2\text{NH}(\text{CH}_3)$ ,



, and







such as in the examples below.









- 20 Pharmaceutically acceptable salts and prodrugs of these compounds are also contemplated.

- 5 Another alteration consists of converting a phenyl moiety to a pyridinyl, furyl, thienyl, or *n*-butyl moiety, such as in the examples below.



- 10 Pharmaceutically acceptable salts and prodrugs of these compounds are also contemplated.

Another alteration consists of adding a substituent comprising from 1 to 3 non-hydrogen atoms to an aromatic or a heteroaromatic ring, as in the examples below.



5    Pharmaceutically acceptable salts and prodrugs of these compounds are also contemplated.

While not intending to limit the scope of the invention, the following are examples of useful compounds

4-{2-[(R)-2-((E)-3-Hydroxy-4-phenyl-but-1-enyl)-6-oxo-  
10    piperidin-1-yl]-ethylsulfanyl}-butyric acid methyl ester, and  
4-{2-[(R)-2-((E)-3-Hydroxy-4-phenyl-but-1-enyl)-6-oxo-  
piperidin-1-yl]-ethylsulfanyl}-butyric acid.

The compounds disclosed herein are useful for the prevention or treatment of glaucoma or ocular hypertension in mammals, or for the  
15    manufacture of a medicament for the treatment of glaucoma or ocular hypertension.

Those skilled in the art will readily understand that for administration or the manufacture of medicaments the compounds disclosed herein can be admixed with pharmaceutically acceptable excipients which per se are well  
20    known in the art. Specifically, a drug to be administered systemically, it may be confectioned as a powder, pill, tablet or the like, or as a solution, emulsion, suspension, aerosol, syrup or elixir suitable for oral or parenteral administration or inhalation.

For solid dosage forms or medicaments, non-toxic solid carriers include,  
25    but are not limited to, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, the polyalkylene glycols, talcum, cellulose, glucose, sucrose and magnesium carbonate. The solid dosage forms may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a  
30    sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated by the technique described in the U.S. Pat. Nos. 4,256,108; 4,166,452; and 4,265,874 to form osmotic therapeutic tablets for control release. Liquid pharmaceutically administrable dosage forms can, for example, comprise  
35    a solution or suspension of one or more of the presently useful compounds and

5 optional pharmaceutical adjutants in a carrier, such as for example, water,  
saline, aqueous dextrose, glycerol, ethanol and the like, to thereby form a  
solution or suspension. If desired, the pharmaceutical composition to be  
administered may also contain minor amounts of nontoxic auxiliary substances  
such as wetting or emulsifying agents, pH buffering agents and the like. Typical  
10 examples of such auxiliary agents are sodium acetate, sorbitan monolaurate,  
triethanolamine, sodium acetate, triethanolamine oleate, etc. Actual methods of  
preparing such dosage forms are known, or will be apparent, to those skilled in  
this art; for example, see Remington's Pharmaceutical Sciences, Mack  
Publishing Company, Easton, Pa., 16th Edition, 1980. The composition of the  
15 formulation to be administered, in any event, contains a quantity of one or more  
of the presently useful compounds in an amount effective to provide the desired  
therapeutic effect.

Parenteral administration is generally characterized by injection, either  
subcutaneously, intramuscularly or intravenously. Injectables can be prepared  
20 in conventional forms, either as liquid solutions or suspensions, solid forms  
suitable for solution or suspension in liquid prior to injection, or as emulsions.  
Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol  
and the like. In addition, if desired, the injectable pharmaceutical compositions  
to be administered may also contain minor amounts of non-toxic auxiliary  
25 substances such as wetting or emulsifying agents, pH buffering agents and the  
like.

The amount of the presently useful compound or compounds  
administered is, of course, dependent on the therapeutic effect or effects desired,  
on the specific mammal being treated, on the severity and nature of the  
30 mammal's condition, on the manner of administration, on the potency and  
pharmacodynamics of the particular compound or compounds employed, and on  
the judgment of the prescribing physician. The therapeutically effective dosage  
of the presently useful compound or compounds is preferably in the range of  
about 0.5 or about 1 to about 100 mg/kg/day.

35 A liquid composition which is intended for topical ophthalmic use is  
formulated such that it can be administered topically to the eye. The comfort

5 should be maximized as much as possible, although sometimes formulation considerations (e.g. drug stability) may necessitate less than optimal comfort. In the case that comfort cannot be maximized, the liquid should be formulated such that the liquid is tolerable to the patient for topical ophthalmic use. Additionally, an ophthalmically acceptable liquid should either be packaged for  
10 single use, or contain a preservative to prevent contamination over multiple uses.

For ophthalmic application, solutions or medicaments are often prepared using a physiological saline solution as a major vehicle. Ophthalmic solutions should preferably be maintained at a comfortable pH with an appropriate buffer  
15 system. The formulations may also contain conventional, pharmaceutically acceptable preservatives, stabilizers and surfactants.

Preservatives that may be used in the pharmaceutical compositions of the present invention include, but are not limited to, benzalkonium chloride, chlorobutanol, thimerosal, phenylmercuric acetate and phenylmercuric nitrate. A  
20 useful surfactant is, for example, Tween 80. Likewise, various useful vehicles may be used in the ophthalmic preparations of the present invention. These vehicles include, but are not limited to, polyvinyl alcohol, povidone, hydroxypropyl methyl cellulose, poloxamers, carboxymethyl cellulose, hydroxyethyl cellulose and purified water.

25 Tonicity adjustors may be added as needed or convenient. They include, but are not limited to, salts, particularly sodium chloride, potassium chloride, mannitol and glycerin, or any other suitable ophthalmically acceptable tonicity adjustor.

Various buffers and means for adjusting pH may be used so long as the  
30 resulting preparation is ophthalmically acceptable. Accordingly, buffers include acetate buffers, citrate buffers, phosphate buffers and borate buffers. Acids or bases may be used to adjust the pH of these formulations as needed.

In a similar vein, an ophthalmically acceptable antioxidant for use in the present invention includes, but is not limited to, sodium metabisulfite, sodium  
35 thiosulfate, acetylcysteine, butylated hydroxyanisole and butylated hydroxytoluene.

5           Other excipient components which may be included in the ophthalmic preparations are chelating agents. A useful chelating agent is edetate disodium, although other chelating agents may also be used in place or in conjunction with it.

The ingredients are usually used in the following amounts:

10

<u>Ingredient</u>	<u>Amount (% w/v)</u>
active ingredient	about 0.001-5
preservative	0-0.10
vehicle	0-40
15   tonicity adjustor	1-10
buffer	0.01-10
pH adjustor	q.s. pH 4.5-7.5
antioxidant	as needed
surfactant	as needed
20   purified water	as needed to make 100%

For topical use, creams, ointments, gels, solutions or suspensions, etc., containing the compound disclosed herein are employed. Topical formulations may generally be comprised of a pharmaceutical carrier, cosolvent, emulsifier, 25 penetration enhancer, preservative system, and emollient.

The actual dose of the active compounds of the present invention depends on the specific compound, and on the condition to be treated; the selection of the appropriate dose is well within the knowledge of the skilled artisan.

30

### Example 1

4-{2-[(R)-2-((E)-3-Hydroxy-4-phenyl-but-1-enyl)-6-oxo-  
piperidin-1-yl]-ethylsulfanyl}-butyric acid methyl ester

Step 1. (R)-2-[2-(3-Methoxycarbonyl-propylsulfanyl)-ethylamino]-hexanedioic acid diethyl ester

35

A mixture of cesium carbonate (2.71 g, 8.32 mmol) and DMF and water (10:1, 20 mL) was stirred at room temperature for 30 min before a solution of (R)-2-

- 5    aminohexanedioic acid diethyl ester (prepared from D- $\alpha$ -  
aminoadipic acid according to Huang, et al., *Synth. Commun.*  
1989, 19, 3485-3496, 1.80 g, 8.28 mmol) in DMF and water  
(10:1, 2 mL) was added via cannula. After 30 min at room  
temperature, potassium iodide (276 mg, 1.66 mmol)
- 10   followed by 4-(2-chloroethylsulfanyl)-butyric acid methyl  
ester (prepared according to PCT 03/007941, 1.63 g, 8.29  
mmol) in DMF and water (10:1, 5 mL) were added. After 23  
h at room temperature, the reaction mixture was heated at  
90 °C. After 2.5 h at 90 °C, the reaction was cooled to room
- 15   temperature and saturated aqueous NaHCO<sub>3</sub> (100 mL) was  
added. The mixture was extracted with EtOAc (3 x 75 mL)  
and the combined extracts were washed with water (2 x 100  
mL) and brine (2 x 100 mL) then dried (Na<sub>2</sub>SO<sub>4</sub>) filtered and  
concentrated in vacuo. Purification of the residue by flash
- 20   column chromatography on silica gel (10% → 40%  
EtOAc/hexane, gradient) two times afforded 893 mg (29%) of  
(R)-2-[2-(3-methoxycarbonyl-propylsulfanyl)-ethylamino]-hexanedioic acid  
diethyl ester.
- 25   Step 2. (R)-1-[2-(3-Methoxycarbonyl-propylsulfanyl)-ethyl]-  
6-oxo-piperidine-2-carboxylic acid ethyl ester  
(R)-2-[2-(3-Methoxycarbonyl-propylsulfanyl)-ethylamino]-hexanedioic acid  
diethyl ester (890 mg, 2.36 mmol), neat, was heated at 100 °C for 18 h. After  
the reaction was cooled to room temperature, TLC and <sup>1</sup>H NMR analysis
- 30   showed no reaction had occurred. The reaction was then heated at 180 °C.  
After 18 h, the reaction mixture was cooled to room temperature.

5 Purification of the residue by flash column chromatography on silica gel (25% EtOAc/hexane → EtOAc, gradient) afforded 163 mg (21%) of (R)-1-[2-(3-methoxycarbonyl-propylsulfanyl)-ethyl]-6-oxo-piperidine-2-carboxylic acid ethyl ester.

10

Step 3. 4-[2-((R)-2-Hydroxymethyl-6-oxo-piperidin-1-yl)-ethylsulfanyl]-butyric acid methyl ester

Lithium borohydride (2.0 M in THF, 0.25 mL, 0.50 mmol) was added slowly to a solution of (R)-1-[2-(3-methoxycarbonyl-propylsulfanyl)-ethyl]-6-oxo-piperidine-2-carboxylic acid ethyl ester (160 mg, 0.48 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1.5 mL) at -40 °C. After 4.5 h at -40 °C, the reaction was quenched by addition of a few drops of aqueous HCl (6 N) until gas evolution ceased. Solid NaHCO<sub>3</sub> was added and the reaction mixture was filtered and concentrated in vacuo. Purification of the residue by flash column chromatography on silica gel (CH<sub>2</sub>Cl<sub>2</sub> → 2% MeOH/CH<sub>2</sub>Cl<sub>2</sub>, gradient) afforded 33 mg (~ 23%) of an inseparable mixture of desired product 4-[2-((R)-2-hydroxymethyl-6-oxo-piperidin-1-yl)-ethylsulfanyl]-butyric acid methyl ester and undesired product (R)-1-[2-(4-hydroxybutylsulfanyl)-ethyl]-6-oxo-piperidine-2-carboxylic acid ethyl ester.

Step 4. 4-[2-((R)-2-Formyl-6-oxo-piperidin-1-yl)-ethylsulfanyl]-butyric acid methyl ester

30



5 1-(3-(Dimethylaminopropyl)-3-ethylcarbodiimide  
hydrochloride (EDCI, 63 mg, 0.33 mmol) and DMSO (31  $\mu$ L,  
0.44 mmol) were added sequentially to a solution of the  
mixture of alcohols from step 3 above (30 mg, ~ 0.10 mmol)  
in benzene (1.5 mL) at 0 °C. After 10 min at 0 °C, pyridinium  
10 trifluoroacetate (23 mg, 0.12 mmol) was added. The reaction was allowed to  
warm to room temperature and then was stirred at room temperature for 2.5 h.  
The solution was decanted from the oily residue and the residue was washed  
with benzene (3 x 2 mL). The combined benzene phases were concentrated in  
vacuo to afford a crude mixture of desired product 4-[2-((R)-2-formyl-6-  
15 oxo-piperidin-1-yl)-ethylsulfanyl]-butyric acid methyl ester  
and undesired (R)-6-oxo-1-[2-(4-oxo-butylsulfanyl)-ethyl]-  
piperidine-2-carboxylic acid ethyl ester.

Step 5. 4-{2-[(R)-2-Oxo-6-((E)-3-oxo-4-phenyl-but-1-enyl)-  
20 piperidin-1-yl]-ethylsulfanyl}-butyric acid methyl ester  
Sodium hydride (60% dispersion in oil, 4.4 mg, 0.11 mmol) was added to a  
solution of dimethyl 2-oxo-3-phenylpropylphosphonate (26.5 mg, 0.11 mmol)  
in THF (0.7 mL) at 0 °C. After 1 h at 0 °C, the mixture of aldehydes from step  
4 above (~ 0.10 mmol) in THF (0.5 mL) was added via cannula. The reaction  
25 was allowed to warm to room temperature. After 18 h at room temperature, the  
reaction was quenched with aqueous acetic acid (50%, 5 mL) and extracted with  
EtOAc (3 x 5 mL). The combined organic phase was washed with brine (10  
mL), dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated in vacuo. Purification of the  
residue by flash column chromatography on silica gel (20% → 60%  
30 EtOAc/CH<sub>2</sub>Cl<sub>2</sub>, gradient) followed by preparative thin layer chromatography  
(silica, 5% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) afforded 6.8 mg (16%) of 4-{2-[(R)-2-oxo-6-  
((E)-3-oxo-4-phenyl-but-1-enyl)-piperidin-1-yl]-  
ethylsulfanyl}-butyric acid methyl ester.

5

Step 6. 4-{2-[(R)-2-((E)-3-Hydroxy-4-phenyl-but-1-enyl)-6-oxo-piperidin-1-yl]-ethylsulfanyl}-butyric acid methyl ester Sodium borohydride (1.0 mg, 0.026 mmol), followed by MeOH (0.1 mL), was added to a solution of 4-{2-[(R)-2-oxo-6-((E)-3-oxo-4-phenyl-but-1-enyl)-piperidin-1-yl]-ethylsulfanyl}-butyric acid methyl ester (6.8 mg, 0.017 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (0.3 mL) at 0 °C. After 15 min at 0 °C, the reaction was quenched with aqueous HCl (0.1 M, 4 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 5 mL). The combined organic phase was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated in vacuo to afford 6.8 mg (99%) of the title compound.

15

#### Example 2

##### 4-{2-[(R)-2-((E)-3-Hydroxy-4-phenyl-but-1-enyl)-6-oxo-piperidin-1-yl]-ethylsulfanyl}-butyric acid

Rabbit liver esterase (134 units/mg, 1 mg) was added to a solution of 4-{2-[(R)-2-((E)-3-hydroxy-4-phenyl-but-1-enyl)-6-oxo-piperidin-1-yl]-ethylsulfanyl}-butyric acid methyl ester (5.2 mg, 0.013 mmol) in acetonitrile (0.2 mL) and pH 7.2 phosphate buffer (3.0 mL). After 24 h, acetonitrile (10 mL) was added and the reaction mixture was concentrated to dryness in vacuo. Purification of the residue by flash column chromatography on silica gel (CH<sub>2</sub>Cl<sub>2</sub> → 5% MeOH/CH<sub>2</sub>Cl<sub>2</sub>, gradient) afforded 4.7 mg (94%) of the title compound.

25

#### Example 3

30

The biological activity of the compounds of Table 1 may be tested using the following procedures.

##### **Radioligand Binding**

5                    **Cells Stably Expressing EP<sub>1</sub>, EP<sub>2</sub>, EP<sub>4</sub> and FP Receptors**

HEK-293 cells stably expressing the human or feline FP receptor, or EP<sub>1</sub>, EP<sub>2</sub>, or EP<sub>4</sub> receptors are washed with TME buffer, scraped from the bottom of the flasks, and homogenized for 30 sec using a Brinkman PT 10/35 polytron. TME buffer is added to achieve a final 40 ml volume in the centrifuge  
10 tubes (the composition of TME is 100 mM TRIS base, 20 mM MgCl<sub>2</sub>, 2M EDTA; 10N HCl is added to achieve a pH of 7.4).

The cell homogenate is centrifuged at 19000 r.p.m. for 20 min at 4° C using a Beckman Ti-60 rotor. The resultant pellet is resuspended in TME buffer to give a final 1 mg/ml protein concentration, as determined by Biorad assay.  
15 Radioligand binding competition assays vs. [<sup>3</sup>H]-17 -phenyl PGF<sub>2α</sub> (5 nM) are performed in a 100μl volume for 60 min. Binding reactions are started by adding plasma membrane fraction. The reaction is terminated by the addition of 4 ml ice-cold TRIS-HCl buffer and rapid filtration through glass fiber GF/B filters using a Brandel cell harvester. The filters are washed 3 times with ice-  
20 cold buffer and oven dried for one hour.

[<sup>3</sup>H]-PGE<sub>2</sub> (specific activity 180 Ci mmol) is used as the radioligand for EP receptors. [<sup>3</sup>H] 17-phenyl PGF<sub>2α</sub> is employed for FP receptor binding studies. Binding studies employing EP<sub>1</sub>, EP<sub>2</sub>, EP<sub>4</sub> and FP receptors are performed in duplicate in at least three separate experiments. A 200μl assay  
25 volume is used. Incubations are for 60 min at 25°C and are terminated by the addition of 4 ml of ice-cold 50 mM TRIS-HCl, followed by rapid filtration through Whatman GF/B filters and three additional 4 ml washes in a cell harvester (Brandel). Competition studies are performed using a final concentration of 5 nM [<sup>3</sup>H]-PGE<sub>2</sub>, or 5 nM [<sup>3</sup>H] 17-phenyl PGF<sub>2α</sub> and non-  
30 specific binding determined with 10<sup>-5</sup>M of unlabeled PGE<sub>2</sub>, or 17-phenyl PGF<sub>2α</sub>, according to receptor subtype studied.

**METHODS FOR FLIPR™ STUDIES**

(a)    **CELL CULTURE**

HEK-293(EBNA) cells, stably expressing one type or subtype of  
35 recombinant human prostaglandin receptors (prostaglandin receptors expressed: hDP/Gqs5; hEP<sub>1</sub>; hEP<sub>2</sub>/Gqs5; hEP<sub>3A</sub>/Gqi5; hEP<sub>4</sub>/Gqs5; hFP; hIP; hTP), are

5 cultured in 100 mm culture dishes in high-glucose DMEM medium containing 10% fetal bovine serum, 2 mM l-glutamine, 250 µg/ml geneticin (G418) and 200 µg/ml hygromycin B as selection markers, and 100 units/ml penicillin G, 100 µg/ml streptomycin and 0.25 µg/ml amphotericin B.

(b) CALCIUM SIGNAL STUDIES ON THE FLIPR™

10 Cells are seeded at a density of  $5 \times 10^4$  cells per well in Biocoat® Poly-D-lysine-coated black-wall, clear-bottom 96-well plates (Becton-Dickinson) and allowed to attach overnight in an incubator at 37 °C. Cells are then washed two times with HBSS-HEPES buffer (Hanks Balanced Salt Solution without bicarbonate and phenol red, 20 mM HEPES, pH 7.4) using a Denley Cellwash  
15 plate washer (LabSystems). After 45 minutes of dye-loading in the dark, using the calcium-sensitive dye Fluo-4 AM at a final concentration of 2 µM, plates are washed four times with HBSS-HEPES buffer to remove excess dye leaving 100 µl in each well. Plates are re-equilibrated to 37 °C for a few minutes.

Cells are excited with an Argon laser at 488 nm, and emission is  
20 measured through a 510-570 nm bandwidth emission filter (FLIPR™, Molecular Devices, Sunnyvale, CA). Drug solution is added in a 50 µl volume to each well to give the desired final concentration. The peak increase in fluorescence intensity is recorded for each well. On each plate, four wells each served as negative (HBSS-HEPES buffer) and positive controls (standard  
25 agonists: BW245C (hDP); PGE<sub>2</sub> (hEP<sub>1</sub>; hEP<sub>2</sub>/Gqs5; hEP<sub>3A</sub>/Gqi5; hEP<sub>4</sub>/Gqs5); PGF<sub>2α</sub> (hFP); carbacyclin (hIP); U-46619 (hTP), depending on receptor). The peak fluorescence change in each drug-containing well is then expressed relative to the controls.

Compounds are tested in a high-throughput (HTS) or concentration-  
30 response (CoRe) format. In the HTS format, forty-four compounds per plate are examined in duplicates at a concentration of  $10^{-5}$  M. To generate concentration-response curves, four compounds per plate are tested in duplicates in a concentration range between  $10^{-5}$  and  $10^{-11}$  M. The duplicate values are averaged. In either, HTS or CoRe format each compound is tested on at least 3  
35 separate plates using cells from different passages to give an  $n \geq 3$ .

5           The results of the activity studies presented in the table will demonstrate that the compounds disclosed herein are have activity characteristic of prostaglandins and are thus useful for the treatment of glaucoma, ocular hypertension, and other diseases or conditions related to prostaglandin activity.


10           The foregoing description details specific methods and compositions that can be employed to practice the present invention, and represents the best mode contemplated. However, it is apparent for one of ordinary skill in the art that further compounds with the desired pharmacological properties can be prepared in an analogous manner, and that the disclosed compounds can also be obtained from different starting compounds via different chemical reactions. Similarly,  
15           different pharmaceutical compositions may be prepared and used with substantially the same result. Thus, however detailed the foregoing may appear in text, it should not be construed as limiting the overall scope hereof; rather, the ambit of the present invention is to be governed only by the lawful construction of the appended claims.

5

CLAIMS

What is claimed is:

1. A compound comprising



10

or a pharmaceutically acceptable salt or a prodrug thereof,


wherein a dashed line indicates the presence or absence of a bond, and an (H) represents a hydrogen atom which is present if required by said bond;

Y is selected from the group consisting of  $\text{CO}_2\text{H}$ ,  $\text{CONMe}_2$ ,  $\text{CONHMe}$ ,

15  $\text{CONHEt}$ ,  $\text{CON}(\text{OCH}_3)\text{CH}_3$ ,  $\text{CONH}_2$ ,  $\text{CON}(\text{CH}_2\text{CH}_2\text{OH})_2$ ,

$\text{CONH}(\text{CH}_2\text{CH}_2\text{OH})$ ,  $\text{CH}_2\text{OH}$ ,  $\text{P}(\text{O})(\text{OH})_2$ ,  $\text{CONHSO}_2\text{CH}_3$ ,  $\text{SO}_2\text{NH}_2$ ,


$\text{SO}_2\text{N}(\text{CH}_3)_2$ ,  $\text{SO}_2\text{NH}(\text{CH}_3)$ ,



R is selected from the group consisting of  $\text{C}_1$ - $\text{C}_4$  alkyl,  $\text{C}_1$ - $\text{C}_4$  alkoxy, halogen,


20  $\text{CO}_2\text{H}$ ,  $\text{OH}$ ,  $\text{COH}$ ,  $\text{COCH}_3$ ,  $\text{COCF}_3$ ,  $\text{NO}_2$ ,  $\text{CN}$ , and  $\text{CF}_3$ .

2. The compound of claim 1 comprising




or a pharmaceutically acceptable salt or a prodrug thereof.

- 5 3. The compound of claim 2 comprising




or a pharmaceutically acceptable salt or a prodrug thereof.

4. The compound of claim 3 consisting of




- 10 5. The compound of claim 1 comprising



or a pharmaceutically acceptable salt or a prodrug thereof.

- 5 6. A compound having an  $\omega$  chain comprising




or a derivative thereof,

wherein a dashed line indicates the presence or absence of a bond, and an (H) represents a hydrogen atom which is present if required by said bond;

- 10 wherein said derivative has a structure as shown above except that an alteration is made to said structure, wherein an alteration consists of

- a. adding, removing, or substituting a non-hydrogen atom of the  $\omega$  chain;
- 15 b. converting a  $\text{CO}_2\text{H}$  to a moiety selected from the group consisting of  $\text{CONMe}_2$ ,  $\text{CONHMe}$ ,  $\text{CONHEt}$ ,  $\text{CON}(\text{OCH}_3)\text{CH}_3$ ,  $\text{CONH}_2$ ,  $\text{CON}(\text{CH}_2\text{CH}_2\text{OH})_2$ ,  $\text{CONH}(\text{CH}_2\text{CH}_2\text{OH})$ ,  $\text{CH}_2\text{OH}$ ,  $\text{P}(\text{O})(\text{OH})_2$ ,  $\text{CONHSO}_2\text{CH}_3$ ,  $\text{SO}_2\text{NH}_2$ ,  $\text{SO}_2\text{N}(\text{CH}_3)_2$ ,  $\text{SO}_2\text{NH}(\text{CH}_3)$ ,




- c. converting a phenyl moiety to a pyridinyl, furyl, thienyl, or *n*-butyl moiety; or
- 20 d. adding a substituent comprising from 1 to 3 non-hydrogen atoms to a phenyl moiety;

or a pharmaceutically acceptable salt or a prodrug thereof.


7. The compound of claim 1 comprising
- 25 4-{2-[(R)-2-((E)-3-Hydroxy-4-phenyl-but-1-enyl)-6-oxo-piperidin-1-yl]-ethylsulfanyl}-butyric acid methyl ester, or



- 5 4-{2-[(R)-2-((E)-3-Hydroxy-4-phenyl-but-1-enyl)-6-oxo-  
piperidin-1-yl]-ethylsulfanyl}-butyric acid,  
or a pharmaceutically acceptable salt or a prodrug thereof.
8. The compound of claim 1 consisting of  
4-{2-[(R)-2-((E)-3-Hydroxy-4-phenyl-but-1-enyl)-6-oxo-  
10 piperidin-1-yl]-ethylsulfanyl}-butyric acid methyl ester, or  
4-{2-[(R)-2-((E)-3-Hydroxy-4-phenyl-but-1-enyl)-6-oxo-  
piperidin-1-yl]-ethylsulfanyl}-butyric acid.
9. A method comprising administering an effective amount of a compound  
to a mammal, said method being effective in treating or preventing glaucoma or  
15 intraocular hypertension, wherein said compound comprises




- or a pharmaceutically acceptable salt or a prodrug thereof,  
wherein a dashed line indicates the presence or absence of a bond, and an (H)  
represents a hydrogen atom which is present if required by said bond;
- 20 Y is selected from the group consisting of CO<sub>2</sub>H, CONMe<sub>2</sub>, CONHMe,  
CONHEt, CON(OCH<sub>3</sub>)CH<sub>3</sub>, CONH<sub>2</sub>, CON(CH<sub>2</sub>CH<sub>2</sub>OH)<sub>2</sub>,  
CONH(CH<sub>2</sub>CH<sub>2</sub>OH), CH<sub>2</sub>OH, P(O)(OH)<sub>2</sub>, CONHSO<sub>2</sub>CH<sub>3</sub>, SO<sub>2</sub>NH<sub>2</sub>,  
SO<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>, SO<sub>2</sub>NH(CH<sub>3</sub>),



- 25 R is selected from the group consisting of C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, halogen,  
CO<sub>2</sub>H, OH, COH, COCH<sub>3</sub>, COCF<sub>3</sub>, NO<sub>2</sub>, CN, and CF<sub>3</sub>.

- 5 10. A liquid composition comprising an effective amount of a compound having an  $\omega$  chain comprising




or a derivative thereof,

wherein a dashed line indicates the presence or absence of a bond, and an (H)

- 10 represents a hydrogen atom which is present if required by said bond;  
wherein said derivative has a structure as shown above except that an alteration is made to said structure, wherein an alteration consists of

- a. adding, removing, or substituting a non-hydrogen atom of the  $\omega$  chain;
- 15 b. converting a  $\text{CO}_2\text{H}$  to a moiety selected from the group consisting of  $\text{CONMe}_2$ ,  $\text{CONHMe}$ ,  $\text{CONHEt}$ ,  $\text{CON}(\text{OCH}_3)\text{CH}_3$ ,  $\text{CONH}_2$ ,  $\text{CON}(\text{CH}_2\text{CH}_2\text{OH})_2$ ,  $\text{CONH}(\text{CH}_2\text{CH}_2\text{OH})$ ,  $\text{CH}_2\text{OH}$ ,  $\text{P}(\text{O})(\text{OH})_2$ ,  $\text{CONHSO}_2\text{CH}_3$ ,  $\text{SO}_2\text{NH}_2$ ,  $\text{SO}_2\text{N}(\text{CH}_3)_2$ ,  $\text{SO}_2\text{NH}(\text{CH}_3)$ ,



- 20 c. converting a phenyl moiety to a pyridinyl, furyl, thienyl, or *n*-butyl moiety; or
- d. adding a substituent comprising from 1 to 3 non-hydrogen atoms to a phenyl moiety;


or a pharmaceutically acceptable salt or a prodrug thereof; and

- 25 wherein said composition is intended for topical ophthalmic use.

5


**ABSTRACT**

A compound comprising



or a pharmaceutically acceptable salt or a prodrug thereof is disclosed herein. Y and R are described in detail herein

10


A compound having an  $\omega$  chain comprising

or a derivative thereof, or a pharmaceutically acceptable salt or a prodrug thereof, is disclosed. Derivatives, salts and prodrugs are identified and described in detail.


15

Methods of treating certain conditions or diseases, and compositions and medicaments related thereto are also contemplated.

**Figure 1**





**Figure 2**



<u>Name</u>	<u>Registration No.</u>
Martin A. Voet	25,208
Robert J. Baran	25,806
Stephen Donovan	33,433
Carlos A. Fisher	36,510
Dean G. Stathakis	54,465

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

<b>FULL NAME OF INVENTOR:</b>			
<b>First Name:</b> <b>DAVID</b>	<b>Initial</b> <b>W.</b>	<b>Last Name</b> <b>OLD</b>	
<b>RESIDENCE &amp; CITIZENSHIP</b>			
<b>City</b> <b>Irvine</b>	<b>State or Foreign Country</b> <b>California</b>	<b>Country Of Citizenship</b> <b>U.S.A.</b>	
<b>POST OFFICE ADDRESS</b>			
<b>Post Office Address</b> <b>13771 Typee Way</b>	<b>City</b> <b>Irvine</b>	<b>State or Country</b> <b>California</b>	<b>Zip Code</b> <b>92620</b>
<b>SIGNATURE OF INVENTOR</b> 		<b>DATE:</b> <b>JUNE 1, 2004</b>	

<b>FULL NAME OF INVENTOR:</b>			
<b>First Name:</b> <b>DANNY</b>	<b>Initial</b> <b>T..</b>	<b>Last Name</b> <b>DINH</b>	
<b>RESIDENCE &amp; CITIZENSHIP</b>			
<b>City</b> <b>Garden Grove</b>	<b>State or Foreign Country</b> <b>California</b>	<b>Country Of Citizenship</b> <b>U.S.A.</b>	
<b>POST OFFICE ADDRESS</b>			
<b>Post Office Address</b> <b>11531 College Avenue</b>	<b>City</b> <b>Garden Grove</b>	<b>State or Country</b> <b>California</b>	<b>Zip Code</b> <b>92840</b>
<b>SIGNATURE OF INVENTOR</b> 		<b>DATE:</b> <b>JUNE 1, 2004</b>	

**APPLICATION DATA SHEET****APPLICATION INFORMATION**

Application Type::	Provisional
Subject Matter::	Utility
Title Line One::	5-THIO-PIPERIDINYL
Title Line Two::	PROSTAGLANDIN E ANALOGS
Title Line Three::	
Attorney Docket Number::	17710 PROV (AP)
Request for Early Publication?::	NO
Request for Non-Publication?::	NO
Suggested Drawing Figure::	
Total Drawing Sheets::	2
Small Entity?::	NO
Petition included?::	NO
Petition Type::	
Licensed US Govt. Agency::	None
Contract or Grant Numbers::	None
Secrecy Order in Parent Appl.?::	NO



**APPLICANT INFORMATION**

**Applicant Authority Type::** Inventor  
**Primary Citizenship Country::** US  
**Status::** Full Capacity  
**Given Name::** DAVID  
**Middle Name::** W.  
**Family Name::** OLD  
**Name Suffix::**  
**City of Residence::** IRVINE  
**State or Province of Residence::** CALIFORNIA  
**Country of Residence::** US  
**Street of mailing address::** 13771 TYPEE WAY  
**City of mailing address::** GARDEN GROVE  
**State or Province of mailing address::** CALIFORNIA  
**Country of mailing address::** US  
**Postal or Zip Code of mailing address::** 92620

**Applicant Authority Type::** Inventor  
**Primary Citizenship Country::** US  
**Status::** Full Capacity  
**Given Name::** DANNY  
**Middle Name::** T.  
**Family Name::** DINH  
**Name Suffix::**  
**City of Residence::** GARDEN GROVE  
**State or Province of Residence::** CALIFORNIA  
**Country of Residence::** US  
**Street of mailing address::** 11531 COLLEGE AVENUE  
**City of mailing address::** GARDEN GROVE  
**State or Province of mailing** CALIFORNIA

**address::**

**Country of mailing address::**

**US**

**Postal or Zip Code of mailing**

**92840**

**address::**

## **CORRESPONDENCE INFORMATION**

### **Correspondence Customer**

**Number::**

**Name Line One::** Brent A. Johnson

**Name Line Two::** Allergan, Inc.

**Street of mailing address::** 2525 Dupont Drive

**City of mailing address::** Irvine

**State or Province of mailing address:** California

**Country of mailing address::** US

**Postal or Zip Code of mailing address::** 92612

**Phone Number::** 714-246-4348

**Fax Number::** 714-246-4249

**E-mail address::** johnson\_brent@allergan.com

### **Representative Information**

<b>Representative Designation::</b>	<b>Registration Number::</b>	<b>Representative Name::</b>
<b>Primary::</b>	51,851	Brent A. Johnson
<b>Associate::</b>	25,208	Martin A. Voet
<b>Associate::</b>	25,806	Robert J. Baran
<b>Associate::</b>	36,510	Carlos A. Fisher
<b>Associate::</b>	33,433	Stephen Donovan
<b>Associate::</b>	54,465	Dean G. Stathakis

**Domestic Priority Information**

Application::    Continuity Type::    Parent Application::    Parent Filing Date::  
This  
Application

**Assignee Information**

Assignee name::	ALLERGAN, INC.
Street of mailing address::	2525 Dupont Drive
City of mailing address::	Irvine
State or Province of mailing address::	California
Country of mailing address::	US
Postal or Zip Code of mailing address::	92612

# **Document made available under the Patent Cooperation Treaty (PCT)**

International application number: PCT/US05/017167

International filing date: 18 May 2005 (18.05.2005)

Document type: Certified copy of priority document

Document details: Country/Office: US  
Number: 60/577,361  
Filing date: 04 June 2004 (04.06.2004)

Date of receipt at the International Bureau: 27 June 2005 (27.06.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse